

Tables

Table 1. Baseline characteristics and clinicopathological stage

		RUNX3 expression		
		High (<i>n</i>)	Low (<i>n</i>)	<i>P</i> value
Age (y) ^a		66.5 ± 10.5	64.7 ± 10.9	0.93
Gender	Male/female	6/4	15/11	0.90
UICC stage	IA	0	2	0.65
	IB	0	0	
	IIA	6	15	
	IIB	4	9	
	III	0	0	
	IV	0	0	
pT	1	0	2	0.37
	2	0	0	
	3	10	24	
	4	0	0	
pN	0	6	17	0.76
	1	4	9	
Histological grade	G1	3	10	0.29

	G2	7	12	
	G3	0	4	
R	0	9	24	0.82
	1	1	2	

^aValues are means \pm SD.

Table 2. Factors predictive of survival after pancreaticoduodenectomy—multivariate analysis

		<i>n</i>	Risk ratio	95% CI	<i>P</i> value
Age (y)	>60/≤60	25/11	1.85	0.66–6.22	0.25
Gender	Male/female	21/15	1.36	0.48–3.79	0.55
UICC stage	IIB + IV/IA + IIA	13/23	3.49	1.38–9.34	0.0083
Histological grade	G3/G1 + G2	4/32	35.4	6.49–211.5	0.0001
Resection margin	R1/R0	3/33	1.14	0.16–4.96	0.87
RUNX3 expression	Low/high	26/10	2.89	1.08–9.38	0.033