

Table 1: Characteristics of patients undergoing the described planned foveal detachment technique for diffuse diabetic macular edema.

No.	Age (years)	Sex	Follow-up (months)	Type	Previous treatment	Preoperative lens status	CRT			BCVA (Snellen)		Continuity of the ellipsoid zone		Continuity of the ELM		Recurrence	Additional treatment for recurrence (no. of treatments)
							Pre	1 week	Final visit	Pre	Final visit	Pre	Final visit	Pre	Final visit		
1	70	M	30	CME	IVB, STTA	Phakic	655	141	213	20/500	20/100	-	-	-	-	+	STTA(3)
2	69	M	27	CME+SRD	IVR, STTA	Pseudophakic	597	375	205	20/29	20/13	+	+	+	+	-	None
3	47	M	26	CME+SRD	IVR, STTA	Pseudophakic	698	340	145	20/29	20/17	+	+	+	+	-	None
4	67	F	25	CME	IVB, STTA	Pseudophakic	558	265	141	20/200	20/200	-	-	-	-	-	None
5	52	M	22	SRS	IVB	Phakic	647	132	124	20/40	20/25	+	+	+	+	-	None
6	53	F	23	CME	IVR	Phakic	508	278	117	20/40	20/33	-	-	-	-	-	None
7	47	M	20	SRS+SRD	IVB, IVR, STTA	Phakic	483	288	116	20/40	20/22	+	+	+	+	-	None
8	68	F	19	SRS	IVR	Pseudophakic	670	444	464	20/50	20/17	+	+	+	+	-	None
9	68	F	18	CME	IVR	Pseudophakic	545	226	129	20/29	20/17	+	+	+	+	-	None
10	66	M	17	CME	IVA, IVR	Phakic	529	225	166	20/50	20/67	+	+	+	+	+	STTA(4)
11	66	F	17	SRS	IVA, IVR	Phakic	484	278	84	20/50	20/33	-	-	-	-	-	None
12	45	F	14	SRS+SRD	IVA	Phakic	930	436	114	20/250	20/200	-	-	-	-	+	STTA(1)
13	70	M	12	CME	IVR	Pseudophakic	771	480	258	20/250	20/100	-	-	-	-	+	STTA(1)
14	64	M	12	SRS	IVR	Pseudophakic	944	133	79	20/400	20/250	-	-	-	-	-	None

CME = cystoid macular edema, SRD = serous retinal detachment, SRS = sponge like retinal swelling, IVB = intravitreal bevacizumab injection, STTA = sub-Tenon's capsule triamcinolone acetate injection, IVR = intravitreal ranibizumab injection, IVA = intravitreal aflibercept injection, CRT = central retinal thickness, BCVA = best-corrected visual acuity, ELM = external limiting membrane