

Table 2. Clinical and radiographic outcomes

	Preoperative	Second-look	P value
Lysholm score	84.8 ± 14.7	97.4 ± 6.4	0.009*
Tegner activity score	4 (4-5)	6 (5-6)	0.017*
IKDC score	63.4 ± 16.3	89.0 ± 8.7	0.002*
KOOS score			
Pain	82.7 ± 18.8	92.5 ± 8.4	0.058
Symptoms	80.6 ± 17.5	89.6 ± 11.7	0.167
Activities of daily life	92.4 ± 13.8	99.4 ± 0.7	0.100
Sport and recreation	63.2 ± 25.5	89.6 ± 9.5	0.005*
Knee related QOL	58.1 ± 20.7	88.0 ± 12.8	0.001*
Total	81.6 ± 16.1	94.4 ± 5.1	0.021*
Side-to-side difference of KT-2000 (mm)	3.6 ± 1.6	0.8 ± 1.6	< 0.001*
Pivot shift test (positive)	17	0	< 0.001*
Kellgren-Lawrence grade (0/1/2/3/4)	18 / 2 / 0 / 0 / 0	18 / 2 / 0 / 0 / 0	1.000

Data of Tegner score are presented as median (interquartile range). Data of Lysholm, International Knee Documentation Committee (IKDC), Knee Injury and Osteoarthritis Outcome Score (KOOS), side-to-side difference of KT-2000 are presented as a mean ± standard deviation. * P < 0.05.

Table 2