

1 Journal of Dermatology Original Article

2

3 **The expression of cell adhesion molecule 1 and its splicing variants in**
4 **Sézary cells and cell lines from cutaneous T-cell lymphoma**

5

6 Mari Yamaguchi¹, Shin Morizane^{1*}, Toshihisa Hamada¹, Tomoko Miyake¹, Makoto

7 Sugaya², Hiroaki Iwata³, Kazuyasu Fujii⁴, Rie Haramoto-Shiratsuki⁵, Yuki Nakagawa¹,

8 Mayumi Miura⁶, Koichi Ohshima⁶, Kazuhiro Morishita⁷, Takahide Takahashi⁸,

9 Masahide Imada^{8,9}, Ken Okada⁸, Jiro Uehara¹⁰, Junko Sowa-Osako¹¹ and Keiji

10 Iwatsuki¹

11

12 ¹Departments of Dermatology, Okayama University Graduate School of Medicine,

13 Dentistry and Pharmaceutical Sciences, Okayama, Japan

14 ²Department of Dermatology, Faculty of Medicine, University of Tokyo, Tokyo, Japan

15 ³Department of Dermatology, Hokkaido University Graduate School of Medicine,

16 Sapporo, Japan

17 ⁴Department of Dermatology, Kagoshima University Graduate School of Medical and

18 Dental Sciences, Kagoshima, Japan

19 ⁵Department of Dermatology, Shimane University Faculty of Medicine, Izumo, Japan

20 ⁶Department of Pathology, Kurume University School of Medicine, Kurume, Japan

21 ⁷Division of Tumor and Cellular Biochemistry, Department of Medical Sciences,

22 Faculty of Medicine, University of Miyazaki, Miyazaki, Japan

23 ⁸Division of Medical Support, Okayama University Hospital, Okayama, Japan

24 ⁹Central Clinical Laboratory, Kawasaki Medical School Hospital, Okayama, Japan

25 ¹⁰Department of Dermatology, Asahikawa Medical University, Asahikawa, Japan

26 ¹¹Department of Dermatology, Osaka City University Graduate School of Medicine,

27 Osaka, Japan

28

29 **Address correspondence to*

30 Keiji Iwatsuki, M.D., Ph.D.

31 Department of Dermatology, Okayama University Graduate School of Medicine,

32 Dentistry and Pharmaceutical Sciences. 2-5-1, Shikata-cho, Kita-ku, Okayama, 700-

33 8558, Japan

34 Phone: +81-86-235-7282, Fax: +81-86-235-7283

35 E-mail: keijiwa@cc.okayama-u.ac.jp

36

37 **Short title**

38 CADM1 expression in Sézary syndrome

39

40 **Abbreviations**

41 cell adhesion molecule-1 (CADM1), tumor suppressor lung cancer-1 (TSLC1), Sézary
42 syndrome (SS), mycosis fungoides (MF), adult T-cell leukemia/lymphoma (ATLL),
43 anaplastic large cell lymphoma (ALCL), C-C chemokine receptor type 4 (CCR4),
44 human T-cell leukemia virus 1 (HTLV-1), peripheral blood mononuclear cell (PBMC),
45 cutaneous T-cell lymphoma (CTCL), diffuse large B-cell lymphoma (DLBCL), enzyme-
46 linked immunosorbent assay (ELISA), reverse transcriptase-polymerase chain reaction
47 (RT-PCR)

48

49

50 Abstract; 242 words (limit: 250 words)

51 Main article; 3087 words (limit: 6,000 words)

52

53 **ABSTRACT**

54 Cell adhesion molecule 1 (CADM1) is aberrantly expressed by T-cell neoplasms such as
55 adult T-cell leukemia/lymphoma (ATLL) and mycosis fungoides (MF). We studied the
56 expression of CADM1 and its splicing variants in Sézary syndrome (SS), MF, other
57 cutaneous T-cell lymphoma (CTCL), and cell lines derived from T- and B-cell
58 lymphomas. Soluble CADM1 was measured in the patients' sera. CADM1+ cells in the
59 blood and skin lesions were examined by flow cytometry and immunostaining,
60 respectively. Soluble CADM1 was measured by ELISA, and the splicing variants of
61 *CADM1* transcripts were determined by reverse transcriptase-polymerase chain
62 reaction, followed by sequencing. As a result, circulating CADM1+ cells were
63 significantly increased in 7 of 10 patients with SS, ranging from 7.9% to 74.5% of the
64 CD3+CD4+ fractions (median; 33.7%) (cut off value; 6.5%). The percentages of
65 CADM1+ cells were usually less than those of circulating Sézary cells. CADM1 was
66 expressed, to various degrees, in 6 of 9 T-cell lines derived from SS, MF, ATLL, and
67 anaplastic large cell lymphoma (ALCL), but negative in B-cell lymphoma-derived cell
68 lines. CADM1+ cells were present in the skin infiltrates of MF, SS, ATLL and ALCL.
69 Serum levels of soluble CADM1 were not significantly elevated in SS/MF. Three major
70 splicing variants of CADM1 expressed by neoplastic T cells contained different

71 combinations of the exons 7, 8, 9 and 11, including a putative oncogenic variant
72 composed of exons 7-8-9-11. In conclusion, CADM1 is frequently expressed in Sézary
73 cells and cell lines from CTCL.

74

75 **Keywords**

76 CADM1, Mycosis fungoides, Sézary syndrome, splicing variant, T-cell lines,

77